CENTRAL INTELLIGENCE AGENCY

REPORT

INFORMATION FROM

FOREIGN DOCUMENTS OR RADIO BROADCASTS

COUNTRY

CD NO.

SUBJECT

Medical Epidemiology

Moscov

DATE OF INFORMATION

HOW **PUBLISHED**

Weakly newspaper

DATE DIST. /4 Jul 1949

1949

WHERE

PUBLISHED

NO. OF PAGES

DATE

PUBLISHED

15 Jun 1949

SUPPLEMENT TO

LANGUAGE

Russian

REPORT NO.

THIS IS UNEVALUATED INFORMATION

SOURCE

Meditsinskiy Rabotnik, No 24, 1949.

MEDICAL GROUP DISCUSSES PROBLEMS OF PROCELLOSIS CONTROL

> Prof V. Troitskiy Corr Mem, Acad Med Sci USSR

Recently the Department of Hygiene, Microbiology, and Epidemiology, Academy of Medical Sciences USSR, convened in special session at Alma-Ata to discuss various aspects of brucellosis and methods for its control. session was well attended by scientific workers of Kazakh 38R, various health officials of the same SSR and guests from various middle Asiatic cities.

The main theme of the session centered around a decree of the Council of Ministers USER, issued early in 1949, which dealt with "Measures for the Control of Brucellosis." Under the program set up by this decree all incidences of brucellosis were to be eliminated by 1950. The USER is the first nation to undertake such a program under government sponsorship.

Much concrete information is necessary towage a successful fight against brucellosis. Scientific and technical personnel must be armed with knowledge concerning the etiology, sources, methods of propagation, development charach teristics, pathogenesis and immunology, as well as methods of prophylaxis and therapy of brucellosis. Therefore, it is necessary to conduct an intensive microbiological, immunological, epidemiological, and clinical study of the disease.

The first report at this memorable session was presented by T. Ye. Boldyrev and I. I. Rogozin, corresponding members of the Academy of Medical Sciences USSA, who described various methods by which state-sponsored institutions can curb the spread of brucellosis. They suggest that the state undertake a comprehensive program to strengthen the sanitation-disinfection network.

CLASSIFICATION MSRB

Sanitized Copy Approved for Release 2011/07/07: CIA-RDP80-00809A000600240231

STAT



Prof P. F. Zdrodovskiy, active member of the Academy, described present-day trends in the study of brucellosis. He revealed that Soviet scientists have established two types of epidemiologically different brucellosis in man: (1) the sporadic type which is transmitted to humans through cows and pigs and (2) the epidemic type which is the result of infection transmitted through goats and sheep. This approach to the study of brucellosis has been most effective and has shown state organizations that every effort must be exerted to eliminate the more dangerous of the two -- the epidenic type.

Academician Ye. N. Pavlovskiy, Dr of Medical Sciences; I. G. Galuzo Director of the Institute of Regional Pathology, Academy of Sciences, Kazakh SSR; I. K. Karakulov; and I. C. Koryakin, Deputy Minister of Health Kazakh SSR; all submitted reports on the present-day status of the epidemiology and epizoetclogy of brucellosis. The reports discussed the role of various animals in the epidemiology of brucellosis.

Pavlovskiy contends that many domestic as well as wild animals are responsible for the spread of brucellosis in nature. Mevertheless, goats and sheep are the chief and primary transmitting agents.

In recent years Soviet scientists have adopted new methods in their studies of brucellosis. They have studied factors governing self-healing, immunity, and the pathogenesis of the dissase. Frof P. F. Zdrodovskiy and his co-workers, P. A. Vershilova and Rh. S. Kotlyarova, described new achievements of USSR science. Studies have shown that self-healing is possible and does occur in brucellosis of the so-called goat and sheep type and the cow type. In the majority of sheep, the cure of brucellosis is accompanied by a complete elimination of causes of the discase. Nevertheless at times the infection recurs, but with less severity. In some cases where sheep have suffered from very acute forms of brucellosis, the infection seems to disappear, but actually remains in the body of the animal in an incubated form. Such studies have been most profitchly since the course of brucellosis in man parallels that of brucellosis in enimals. Date, such as this, has brought about a complete change in the pathogenetical conceptions of this disease.

Immunity from brucellosis can now be studied from a new view point because of knowledge concerning the dynamics of infection and common characteristics of the self-healing of the man or beast afflicted with brucellosis. Soviet scientists have been able to develop not only nonsterile v. .cine for use during infection but also a postinfection sterile vaccine. The latter is used to prevent re-infection of the patient. In addition to problems revolving around the possibilities of self-healing and postinfection immunization, there is also the problem of a live vaccine to prevent infection from the disease. P. A. Vershilowa, senior scientific collaborator, submitted a report which dealt with the possibility of developing just such a live vaccine. Experiments have shown that it was possible to obtain a dry, live avirulent brucellosis vaccine, which has great practical value.

Many of the reports submitted dealt with the clinical aspects and treatment of hymnellosis. The most important was submitted by Prof G. P. Rudneva, Corresponding Member of the Academy of Medical Sciences USSR. Prof O. D. Sokolova-Ponomars va, Corresponding Member of the Academy, reported on the clinical aspects of trucellosis in children. Docent M. Kh. Farizova reported on the clinical aspects of nerve disorders brought about by brucellosis infection. N. D. Beklemishev, Candidate of Medical Sciences, described the significance of allergy in the treatment of brucellosis. Prof P. P. Ochkur, Chief of the Chair of Pathology, Kazakh Medical Institute discussed the pathological morphology of brucellosis in man.





STAT

All the reports showed the great interest and achievements of Soviet scientists in the field of brucellosis and its cure. Along this line Prof G. T. Rudnev's report introduced a note of optimism for all who might be suffering from this disease. It is now possible to give prompt aid and in many cases effect a cure in brucellosis 'nfection. An effective cure consists of close correlation between immunobiological, chemotherapeutical and physicotherapeutical methods. In recent years the effectiveness of vaccination has become very evident.

Others who submitted reports were M. N. Solov'yev, P. G. Sergiyev, active members of the Academy; S. N. Puchkovskiy, A. Ya. Alymov, P. A. Petrishcheva, corresponding members of the Academy; and various representatives of sowkhozes, kolkhozes, and rural regions of Fazakh SSR.

- E N D -